



→ **OTHER**
e.g. Lice, scrapie,
mycotic dermatitis etc

SHEEP SCAB CONFIRMED

DIP

NOT SHOWER OR JET

INJECT

(DOSE FOR HEAVIEST IN GROUP)

OP Organophosphate Coopers Ectoforce Paracide Plus Osmonds Gold Fleece	SP Long acting High-cis cypermethrin Ectofleece Auriplak	SP Short acting High-cis cypermethrin Robust
1 treatment gives up to 4 weeks protection		No protection against re-infection. Redip after 14 days
35	12	18 after 2 nd dip
BROAD SPECTRUM Kills scab mites, lice and other ectoparasites		

PRODUCTS

PROTOCOL & PROTECTION

WITHDRAWAL PERIOD (DAYS)

FEATURES

MOXIDECTIN Cydectin Zermex	DORAMECTIN Dectomax	IVERMECTIN Ivomec Depidex Virbamec Noromect Paramectin
2 injections SUBCUTANEOUS to treat scab 10 days apart. Must keep treated sheep isolated for 12 days after treatment Up to 28 days protection	1 injection. INTRAMUSCULAR Must keep treated sheep isolated on clean area* for 14 days after treatment	2 injections SUBCUTANEOUS 7 days apart. Must keep treated sheep isolated on clean area* for 7 days after treatment
* clean = no sheep in previous 21 days		
70 after 2 nd injection	70	42 after 2 nd injection
SCAB ONLY (and susceptible worms)		

OTHER ACTIONS

1. Establish where the scab came from → If purchased or incoming sheep review quarantine protocols.
→ Neighbours (see below)
2. Contact neighbours with sheep in adjacent fields to warn them and/or suggest they treat at the same time as you for maximum effect and protection.
3. Check common fence-lines for gaps/shared rubbing areas. Consider double fencing any in contact with high risk neighbours.

ALWAYS READ THE MANUFACTURER'S INSTRUCTIONS BEFORE USE